

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
7 September 2001 (07.09.2001)

PCT

(10) International Publication Number
WO 01/64660 A1(51) International Patent Classification: **C07D 265/00**,
A61K 31/555, C07F 5/02**B.** [US/US]; 11977 Neihaus Lane, St. Louis, MO 63146 (US).

(21) International Application Number: PCT/US01/06394

(74) Agent: LIMPUS, Lawrence, L.; 675 McDonnell Boulevard, P.O. Box 5840, St. Louis, MO 63134 (US).

(22) International Filing Date: 28 February 2001 (28.02.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
09/517,252 2 March 2000 (02.03.2000) US(71) Applicant (for all designated States except US):
MALLINCKRODT INC. [US/US]; 675 McDonnell Boulevard, P.O. Box 5840, St. Louis, MO 63134 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **RAJAGOPALAN, Raghavan** [US/US]; 13031 Vinson Court, Maryland Heights, MO 63043 (US). **ACHILEFU, Samuel, I.** [NG/US]; 3424 San Sevilla Court, St. Louis, MO 63044 (US). **BUGAJ, Joseph, E.** [US/US]; 2916 Kettering Drive, St. Charles, MO 63303 (US). **DORSHOW, Richard,**

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: NOVEL QUINOLINE LIGANDS AND METAL COMPLEXES FOR DIAGNOSIS AND THERAPY

(57) Abstract: The present invention relates to novel ligands for forming metal complexes that absorb or fluoresce in the visible or near-infrared (NIR) region of the electromagnetic spectrum, new complexes incorporating such ligands, process for preparing such complexes, and methods of imaging or therapy using such agents. More particularly, the present invention specifically pertains to novel metal complexes derived from quinoline based heterocyclic N_2O_3 , N_2O_4 , N_2O_5 , N_2O_6 and N_2O_7 ligands, and are useful as general imaging, diagnostic, or therapeutic agents employing optical, nuclear medicine, or magnetic resonance procedures.

NOVEL QUINOLINE LIGANDS AND METAL COMPLEXES FOR DIAGNOSIS AND THERAPY

FIELD OF THE INVENTION

5

The present invention relates to diagnosis and therapy within the field of biomedical optics. More particularly, the invention relates to novel ligands for forming metal complexes that absorb or fluoresce in the visible or near-infrared (NIR) region of the electromagnetic spectrum, new complexes
10 incorporating such ligands, process for preparing such complexes, and methods of imaging or therapy using such agents.

BACKGROUND OF THE INVENTION

15

The field of biomedical optics is growing rapidly due to distinct advantages over other imaging modalities such as X-ray CT, MRI, nuclear medicine, or ultrasound (J.C. Hebden and D.T. Delpy. Diagnostic Imaging with Light, *The British Journal of Radiology*, **1997**, 70, S206-S214; G. Freiherr. The Light Stuff: Optical Imaging in Medical Diagnostics, *Medical
20 Devices & Diagnostic Industry*, **1998**, 40-46). Compounds absorbing or emitting in the visible or NIR region of electromagnetic spectrum are potentially useful for tomographic imaging, endoscopic examination, photodynamic therapy, optoacoustic imaging, and sonofluorescence techniques. Furthermore, compounds absorbing or emitting in the
25 appropriate visible region can be used to generate singlet oxygen and have been shown to be effective for photodynamic therapy of certain types of tumors.

Metal ions continue to play a major role in diagnostic and therapeutic medicine. For example, radionuclide metal complexes derived from both

- transition and lanthanide elements are being used extensively in diagnostic and therapeutic nuclear medicine procedures, paramagnetic complexes are being used extensively in magnetic resonance imaging procedures, and platinum complexes have long been used as cancer chemotherapeutic agents.
- 5 Recently, metal complexes that absorb or emit in the visible or near-infrared (NIR) region have made a significant impact in the field of biomedical optics and have a great potential for photodiagnostic and phototherapeutic applications (J.N. Demas and B.A. DeGraff. Design and Applications of Highly Luminescent Transition Metal Complexes, *Analytical Chemistry*, 1991, 63, 829-837; M.P. Houline et al. Spectroscopic Characterization and Tissue Imaging Using Site-Selective Polyazacyclic Terbium (III) Chelates, *Applied Spectroscopy*, 1996, 50(10), 1221-1228; J.R. Lakowicz et al. Development of Long-Lifetime Metal-Ligand Probes for Biophysics and Cellular Imaging, *Journal of Fluorescence*, 1997, 7, 17-25; F.J. Steemers et al. Near-Infrared Luminescence of Yb^{3+} , Nd^{3+} , and Er^{3+} Azatriphenylene Complexes, *Tetrahedron Letters*, 1998, 39, 7583-7586; G.E. Keifer and D.J. Bornhop. Fluorescent Chelates as Visual Tissue Specific Imaging Agents, *US Patent*, 1999, 5922867). Examples of suitable metal ions for optical applications include Cr(III), Os(II), Ru(II), Ni(II), Eu(III), Tb(III), Lu(III), 15 Yb(III), Er(III), and Nd(III). Eu(III), and Tb(III) are particularly preferred because of favorable absorption and emission properties in visible and NIR regions.

- The key requirements for design of novel metal complexes for optical diagnostic and therapeutic application are: (a) strong absorption and 25 emission in the visible or NIR region; (b) high thermodynamic, kinetic, and photo stability; (c) low toxicity; (d) water solubility; and (e) conjugation capability for targeted delivery to particular tissues or organs. Free metal ions are generally quite toxic; they need to be administered in the form of complexes with complexing agents (ligands) in order to deliver them to

- specific organs and to alleviate toxicity. Electronic property, toxicity, stability, and tissue specificity are greatly affected by the nature of the complexing agents (ligands). Various physicochemical and pharmacokinetic factors have to be considered in order to render the metal complex safe and effective. Electronic requirements for enabling the metal ion (transition and lanthanide) to absorb or emit in the visible or NIR region are well established and essentially involve incorporation of metal into highly polarizable π -electron rich, multidentate ligand systems. Energy transfer from aromatic donors (referred to as "antennae") to the lanthanide metal ion directly bounded to the donor group results in large increase in lanthanide fluorescence (S.I. Weissman, *Journal of Chemical Physics*, **1942**, 10, 214; B. Alpha et al. Energy Transfer Luminescence of Europium (III) and Terbium (III) Cryptates of Macrobicyclic Polypyridine Ligands, *Angewandte Chemie International Edition in English*, **1987**, 26(3), 266-267; J.B. Lamture et al. Luminescence Properties of Terbium (III) Complexes with 4-Substituted Dipicolinic Acid Analogues, *Inorganic Chemistry*, **1995**, 34, 864-869). In contrast, simple lanthanide metal salts or lanthanide ions coordinated to polyaminocarboxylate ligands wherein the aromatic donors are not directly attached exhibit very weak fluorescence in aqueous media (A. Abusaleh and C.F. Meares. *Photochemistry and Photobiology*, **1984**, 39, 763-769). Long-wavelength fluorescence of transition metal complexes generally occurs via metal-to-ligand charge transfer (MCLT) interactions (Z. Murtaza and J.R. Lakowicz. Long-lifetime and Long-wavelength Osmium (II) Metal Complexes Containing Polypyridine Ligands. Excellent Red Fluorescent Dyes for Biophysics and for Sensors, *SPIE*, **1999**, 3602, 309-315).

Toxicity of metal complexes is greatly affected by the nature of the ligands. Since *in vivo* release of free metal ions from the complex is a major cause of toxicity, thermodynamic and kinetic stability are critical requirements for the design of novel ligands. The thermodynamic stability

constant indicates the affinity of totally unprotonated ligand for a metal ion. The conditional stability constant indicates the stability of the complex under physiological pH. Ion selectivity of the ligand toward the desired metal ion over other endogenous metal ions such as zinc, iron, magnesium, and calcium. determines the rate of release of the metal ion into the vascular or extracellular space. The released metal ion is capable of crossing the blood-brain barrier and thereby perturbing the neurophysiology. Therefore, *in vivo* reaction kinetics are also a major factor in the design of stable complexes and complexes with structural features that make *in vivo* transmetallation reactions proceed much slower than the biological clearance of the intact metal complexes would be predicted to have low toxicities (W. Cacheris et al., *Magnetic Resonance Imaging*, **1990**, 8, 467; Oksendal et al., *Journal of Magnetic Resonance Imaging*, **1993**, 3, 157). Thus, a need continues to exist for new and structurally diverse metal complexes for use as imaging, diagnostic, or therapeutic agents employing optical procedures.


SUMMARY OF THE INVENTION

Thermodynamically and kinetically stable metal complexes can be achieved with a proper choice of ligands systems. Transition metal ions generally require soft donors such as thiols and phosphines, whereas lanthanide ions require hard donors such as carboxylates or amines. However, unsaturated heterocyclic bases such as pyridines, imidazoles, and the like are excellent coordinators to both types of metal ions. Numerous pyridine, quinoline, and imidazole based metal complexes have been prepared and many of them have been conjugated to bioactive carriers such as immunoglobulins (R. Rajagopalan et al. Preparation, Characterization, and Biological Evaluation of Technetium (V) and Rhenium (V) Complexes of Novel Heterocyclic Tetradentate N₃S Ligands, *Bioconjugate Chemistry*,

1997, 8, 407-415; J.B. Lamture and T.G. Wensel. A Novel Reagent for Labeling Macromolecules with Intensely Luminescent Lanthanide Complexes, *Tetrahedron Letters*, **1993**, 34(26), 4141-4144). The present invention specifically pertains to novel quinoline based heterocyclic N_2O_3 , N_3O_3 , N_3O_4 , and N_2OS ligands that are suitable for complexing metal ions, and are useful as general imaging, diagnostic, or therapeutic agents employing optical, nuclear medicine, or magnetic resonance procedures. The principal advantages of this invention are: (a) the metal ion is directly bounded to the "antenna" portion of the molecule, and (b) the entire complex is rigid. Both of these factors are expected to contribute to significant enhancement of absorption and luminescence properties compared to those metal complexes where the antenna is either located remote from the metal ion or has considerable degrees of freedom.

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides new and structurally diverse compositions comprising complexing agents (ligands) of the general Formula 1,



Formula 1

wherein R¹ to R⁵ may be the same or different and are selected from the group consisting of hydrogen, C1-C10 alkyl, -OH, C1-C10 polyhydroxyalkyl, C1-C10 alkoxy, C1-C10 alkoxyalkyl, -SO₃H, -(CH₂)_m-CO₂H and -NR⁶R⁷; R⁶ and R⁷ may be the same or different and are selected from the group consisting of


hydrogen, C1-C10 alkyl, C1-C10 aryl, and C1-C10 polyhydroxyalkyl; m ranges from 0 to 10; A¹ is selected from the group consisting of -OH, -CO₂H, -N(R⁸)(R⁹), -SPg-CONHR¹⁰ and -HNCOR¹¹; R⁸ and R⁹ may be the same or different and are selected from the group consisting of hydrogen, C1-C10 alkyl, C1-C10 aryl, C1-C10 polyhydroxyalkyl, -(CH₂)_mCO₂H and -(CH₂)₂-N(CH₂CO₂H)₂; R¹⁰ is selected from the group consisting of hydrogen, C1-C10 alkyl, C1-C10 aryl, C1-C10 polyhydroxyalkyl and -(CH₂)₂-SPg; R¹¹ is selected from the group consisting of hydrogen, C1-C10 alkyl, C1-C10 aryl, C1-C10 polyhydroxyalkyl and -CH(R¹²)-SPg; R¹² is selected from the group consisting of hydrogen, C1-C10 alkyl, C1-C10 aryl, C1-C10 polyhydroxyalkyl, -(CH₂)_mCO₂H and -(CH₂)_nNH₂; n varies from 1 to 10; B¹ is selected from the group consisting of -CHR¹³ and -CH(R¹⁴)(R¹⁵); R¹³ to R¹⁵ may be the same or different and are defined in the same manner as R¹²; C¹ is selected from the group consisting of hydrogen, C1-C10 alkyl, C1-C10 aryl, C1-C10 polyhydroxyalkyl, C1-C10 alkoxy, C1-C10 alkoxyalkyl, -(CH₂)_mCO₂H, -CH₂CH₂-N(CH₂CO₂H)₂, -(CH₂)₂-SPg and -COCH(R¹⁶)-SPg; R¹⁶ is defined in the same manner as R¹²; D¹ is selected from the group consisting of hydrogen, C1-C10 alkyl, C1-C10 aryl, hydroxyl, C1-C10 polyhydroxyalkyl, C1-C10 alkoxy, C1-C10 alkoxyalkyl, -(CH₂)_mCO₂H, and -CH₂CH₂-N(CH₂CO₂H)₂; and Pg may be hydrogen or a protecting group selected from the group consisting of t-butyl, benzyl, benzoyl, 2,4-dimethoxybenzyl, trityl, tetrahydropyranyl, C1-C10 acyl, C1-C10 alkoxyalkyl, and C1-C10 alkoxycarbonyl.

In a preferred embodiment, ligands according to the present invention have the general formula of Formula 1 above wherein R¹ to R⁵ are selected from the group consisting of hydrogen, -OH, C1-C10 alkoxy, -(CH₂)_m-CO₂H, and -N(R⁶)(R⁷); A¹ is selected from the group consisting of -OH, -N(R⁸)(R⁹), and -HNCOR¹¹; B¹ is -CHR¹³; C¹ is selected from the group consisting of hydrogen, C1-C10 alkyl, -(CH₂)_mCO₂H, -CH₂CH₂-

$N(CH_2CO_2H)_2$, and $-COCH(R^{16})-SPg$; D^1 is selected from the group consisting of hydrogen, C1-C10 alkyl, C1-C10 polyhydroxyalkyl, $-(CH_2)_m-CO_2H$, and $-CH_2CH_2-N(CH_2CO_2H)_2$; and Pg is selected from the group consisting of C1-C10 acyl, tetrahydropyranyl, C1-C10 alkoxyalkyl, and C1-
 5 C10 alkoxycarbonyl.

In a further preferred embodiment, ligands according to the present invention have the general formula of Formula 1 above wherein R^1 to R^5 are hydrogens; A^1 is $-OH$ or $-N(CH_2CO_2H)_2$; B^1 is $-CH_2-$; C^1 is selected from the group consisting of $-CH_2-CO_2H$, $-CH_2CH_2-N(CH_2CO_2H)_2$, and $-COCH_2-SPg$;
 10 D^1 is selected from the group consisting of hydrogen, C1-C10 alkyl, and $-CH_2-CO_2H$; and Pg is selected from the group consisting of benzoyl, tetrahydropyranyl, and methoxycarbonyl.

The present invention also provides structurally diverse compositions comprising metal complexes of the general formula of Formula 2 formed by
 15 coordination of an appropriate metal ion to the ligands derived from Formula 1 shown above,



Formula 2

wherein R^{17} to R^{21} may be the same or different and are defined in the same manner as R^1 ; A^2 is selected from the group consisting of $-O-$, $-CO_2-$, $-N(R^8)(R^9)$, $-SPg-CON(R^{10})$, and $-NCOR^{11}$; R^8 and R^9 may be the same or
 20 different and are selected from the group consisting of hydrogen, C1-C10 alkyl, C1-C10 aryl, C1-C10 polyhydroxyalkyl, $-(CH_2)_mCO_2-$, and $-(CH_2)_2-N(CH_2CO_2-)_2$; R^{10} is selected from the group consisting of hydrogen, C1-C10 alkyl, C1-C10 aryl, C1-C10 polyhydroxyalkyl, and $-(CH_2)_2-S-$; R^{11} is selected

from the group consisting of hydrogen, C1-C10 alkyl, C1-C10 aryl, C1-C10 polyhydroxyalkyl, and $-\text{CH}(\text{R}^{12})-\text{S}^-$; R^{12} is selected from the group consisting of hydrogen, C1-C10 alkyl, C1-C10 aryl, C1-C10 polyhydroxyalkyl, $-(\text{CH}_2)_m\text{CO}_2^-$ and $-(\text{CH}_2)_n\text{NH}_2$; n varies from 1 to 10; B^2 is selected from the group consisting of $-\text{CHR}^{13}$ and $-\text{CH}(\text{R}^{14})\text{CH}(\text{R}^{15})$; R^{13} to R^{15} may be the same or different and are defined in the same manner as R^{12} ; C^2 is selected from the group consisting of hydrogen, C1-C10 alkyl, C1-C10 aryl, C1-C10 polyhydroxyalkyl, C1-C10 alkoxy, C1-C10 alkoxyalkyl, $-(\text{CH}_2)_m\text{CO}_2^-$, $-\text{CH}_2\text{CH}_2-\text{N}(\text{CH}_2\text{CO}_2^-)_2$, $-(\text{CH}_2)_2-\text{S}^-$, and $-\text{COCH}(\text{R}^{16})-\text{S}^-$; R^{16} is defined in the same manner as R^{12} ; D^2 is selected from the group consisting of hydrogen, C1-C10 alkyl, C1-C10 aryl, hydroxyl, C1-C10 polyhydroxyalkyl, C1-C10 alkoxy, C1-C10 alkoxyalkyl, $-(\text{CH}_2)_m\text{CO}_2^-$, and $-\text{CH}_2\text{CH}_2-\text{N}(\text{CH}_2\text{CO}_2^-)_2$; and M is a metal ion having an atomic number of 24 to 31, 42 to 49, 58-71, or 74-83.

In a preferred embodiment, the complexes according to the present invention have the general formula of Formula 2 above wherein R^{17} to R^{21} are selected from the group consisting of hydrogen, $-\text{O}^-$, C1-C10 alkoxy, $-(\text{CH}_2)_m-\text{CO}_2^-$, and $-\text{N}(\text{R}^6)(\text{R}^7)$; A^2 is selected from the group consisting of $-\text{O}^-$, $-\text{N}(\text{R}^8)(\text{R}^9)$, and $-\text{HNCOR}^{11}$; B^2 is $-\text{CHR}^{13}$; C^2 is selected from the group consisting of hydrogen, C1-C10 alkyl, $-(\text{CH}_2)_m\text{CO}_2^-$, $-\text{CH}_2\text{CH}_2-\text{N}(\text{CH}_2\text{CO}_2^-)_2$, and $-\text{COCH}(\text{R}^{16})-\text{S}^-$; D^2 is selected from the group consisting of hydrogen, C1-C10 alkyl, C1-C10 polyhydroxyalkyl, $-(\text{CH}_2)_m-\text{CO}_2^-$, and $-\text{CH}_2\text{CH}_2-\text{N}(\text{CH}_2\text{CO}_2^-)_2$; and M is a metal ion having an atomic number of 24 to 28, 31, 42 to 45, 49, 62, 65, 71, 75, 76, or 83.

In a further preferred embodiment, the complexes according to the present invention have the general formula of Formula 2 above wherein R^{17} to R^{21} are hydrogens; A^2 is $-\text{O}^-$ or $-\text{N}(\text{CH}_2\text{CO}_2^-)_2$; B^2 is $-\text{CH}_2-$; C^2 is selected from the group consisting of $-\text{CH}_2-\text{CO}_2^-$, $-\text{CH}_2\text{CH}_2-\text{N}(\text{CH}_2\text{CO}_2^-)_2$, and $-\text{COCH}_2-\text{S}^-$; D^2 is selected from the group consisting of hydrogen, C1-C10

alkyl, and $-\text{CH}_2\text{-CO}_2^-$; and M is a metal ion having an atomic number of 24-26, 28, 31, 43, 44, 49, 62-65, 71, 75, or 76.

The compositions of the invention are suitable for use with a variety of other modalities including X-rays, magnetic resonance, and radiographic
5 imaging. Electron donating and electron releasing groups at various positions in the ligands of Formula 1 and the metal complexes Formula 2 provide an opportunity to alter the absorption and emission properties of the molecule thereby enhancing the optical utility of these molecules. Also, these additional functionalities afford the capability of conjugation to biomolecules
10 and synthetic polymers for selective delivery to various organs or tissues of interest. The term biomolecule refers to all natural and synthetic molecules that play a role in biological systems. Biomolecules include hormones, amino acids, peptides, peptidomimetics, proteins, nucleosides, nucleotides, nucleic acids, carbohydrates, lipids, albumins, mono- and polyclonal
15 antibodies, receptor molecules, receptor binding molecules, synthetic polymers, and aptamers. Specific examples of biomolecules include inulins, prostaglandins, growth factors, growth factor inhibitors like somatostatin, liposomes, and nucleic acid probes. Example of synthetic polymers include polylysine, polyaspartic acid, polyarginine, aborols, dendrimers, and
20 cyclodextrins. The advantages of biomolecules include enhance tissue targeting through specificity and delivery. The specific targeting of effector molecules to a particular tissue, such as tumor, using antibodies is well known in the art (see Halpern et al., Diagnostic Imaging, 1983, 40). Coupling of bifunctional ligands and complexes to biomolecules can be
25 accomplished by several known methods (see Hnatowich et al., Science, 1983, **220**, 613).

The complexes of the present invention may vary widely depending on the contemplated application. For diagnostic imaging of areas of lesion, fluorescent compounds absorbing and emitting in the near infrared (NIR)

region, i.e. 650-900 nm, are desirable. For monitoring blood clearance or for endoscopic examination of lesions, dyes absorbing and emitting in the region of 350-950 nm, preferably 600-900 nm, are desirable. Similarly, the carrier molecules may also vary widely. For blood persistent agents, albumin or

5 methylated serum albumin is preferable. For renal function measurements, polysaccharides or anionic polypeptides are desirable. For endoscopic examination of lesions, antibodies, peptides, or carbohydrates directed against specific cell surface markers are preferred.

Diagnostic compositions comprising the compounds of the invention

10 are also provided. Methods of performing diagnostic procedures with compositions of the invention are also disclosed. The method comprises administering an effective amount of a composition of the invention contained in a pharmaceutically acceptable formulation to a patient either systemically or locally to the organ or tissue to be studied. It is believed that


15 the novel compositions of the present invention have broad clinical utility, which includes, but is not limited to, diagnostic imaging of tumors, of inflammation (both sterile and bacterial), and of impaired vasculature; laser guided endoscopic examination of sites of lesion; and photodynamic and chemotherapy of tumors or infection.

20 The novel compositions of this invention can be formulated into diagnostic or therapeutic compositions for enteral, parenteral, or oral administration. These compositions contain an effective amount of the metal complexes along with conventional pharmaceutical carriers and excipients appropriate for the type of administration contemplated. These compositions

25 may also include stabilizing agents selected from the class consisting of mono- or polycarboxylic acids, mono- or polyamines, mono- or polynucleotides, mono or polysaccharides, amino acids, and peptides. For example, parenteral administration advantageously contains a sterile aqueous solution or suspension of the complexes whose concentration ranges from

about 1 nM to about 0.5 M. Preferred parenteral formulations have a concentration of 1 μ M to 10 mM. Such solutions also may contain pharmaceutically acceptable buffers, emulsifiers, surfactants, and, optionally, electrolytes such as sodium chloride. Concentrations of the metal complexes of this invention in formulations for enteral administration may vary widely as is well-known in the art. In general, such formulations are liquids which include an effective amount of the complexes in aqueous solution or suspension. Such enteral composition may optionally include buffers, surfactants, emulsifiers, thixotropic agents, and the like. Compositions for oral administration may also contain flavoring agents and other ingredients for enhancing their organoleptic qualities. The diagnostic compositions are administered in doses effective to achieve the desired diagnostic or therapeutic objective. Such doses may vary widely depending upon the particular complex employed, the organs or tissues to be examined, the equipment employed in the clinical procedure, and the like.


The following examples illustrate specific embodiments of this invention. As would be apparent to skilled artisans, various changes and modifications are possible and are contemplated within the scope of the invention described.

Example 1. Preparation of the quinoline ligand (Formula 3).

Formula 3


5 A mixture of 2-aminomethyl-8-hydroxyquinoline hydrochloride (Chem-Master International) (2.1 g, 10 mmol), t-butyl bromoacetate (3.9 g, 20 mmol), diisopropylethylamine (3.9 g, 30 mmol), and sodium iodide (0.15 g, 1 mmol) in dimethoxyethane (20 mL) was heated under reflux for 2 hours. The reaction mixture was poured onto water and extracted with ethyl acetate
 10 (3 x 30 mL). The combined organic layers were washed with water (3 x 50 mL), dried (Na₂SO₄), filtered, and the filtrate evaporated in vacuo to furnish 2.2 g of the diester as a red gum. ¹H-NMR (CDCl₃) δ 8.17 (s, 1H), 7.60 (d, 1H), 7.40 (d, 1H), 7.35 (m, 2H), 7.15 (d, 1H), 4.18 (s, 2H), 3.50 (s, 4H); ¹³C-NMR (CDCl₃) δ 165.7, 152.7, 147.2, 137.8, 122.4, 117.4, 112.8, 105.0, 76.4,
 15 55.0, 50.9, 23.3.

The diester obtained above was treated with 96% formic acid (20 mL) and kept at ambient temperature for 24 hours. Excess formic acid was removed by evaporation in vacuo and the brown residue was treated with water (25 mL), and filtered hot (gravity filtration). Upon cooling, the product
 20 crystallized as a red solid which was filtered and dried to furnish 1.1 g of the ligand of Formula 3. ¹H-NMR (DMSO-d₆) δ 9.45 (broad, 2H), 8.15 (d, 1H), 7.77 (d, 1H), 7.28 (m, 2H), 7.10 (d, 1H), 4.15 (s, 2H), 3.50 (s, 4H); ¹³C-NMR (DMSO-d₆) δ 172.5, 158.1, 152.8, 137.4, 136.4, 127.9, 127.0, 121.5, 117.6, 111.3, 59.7, 54.5; electrospray mass spectrum, m/z = 291 (M+H).

Example 2. Preparation of the chromium complex (Formula 4).

Formula 4

- A solution of the ligand of Formula 3 (145 mg, 0.5 mmol) and
 5 chromium acetylacetonate (175 mg, 0.5 mmol) in dimethylformamide (2 mL)
 was treated with two drops of water and two drops of 96% formic acid and
 the entire mixture was heated at 100-120 °C for 24 hours. After cooling the
 reaction mixture to ambient temperature, the solution was poured onto ethyl
 ether. The brown precipitate was collected by filtration, washed with ether,
 10 dried, and recrystallized from propanol to give the chromium complex of
 Formula 4 as a brown solid.


Example 3. Hypothetical preparation of the iron complex (Formula 5).

Formula 5

- A solution of the ligand of Formula 3 (145 mg, 0.5 mmol) and iron
 acetylacetonate (177 mg, 0.5 mmol) in dimethylformamide (2 mL) is treated
 with two drops of water and the entire mixture is heated at 100-120 °C for 16
 hours. After cooling the reaction mixture to ambient temperature, the
 20 solution is poured onto ethyl ether. The precipitate is collected by filtration


and is purified by either recrystallization or C-18 reverse phase chromatography to give the iron complex of Formula 5.

Example 4. Preparation of ruthenium complex (Formula 6).



Formula 6

A solution of the ligand of Formula 3 (145 mg, 0.5 mmol) and ruthenium acetylacetonate (199 mg, 0.5 mmol) in dimethylformamide (2 mL) was treated with two drops of water and the entire mixture was heated at 100-120 °C for 16 hours. After cooling the reaction mixture to ambient temperature, the solution was poured onto ethyl ether. The brown precipitate was collected by filtration, washed with ether, dried, and recrystallized from propanol to give the ruthenium complex of Formula 6 as a black solid.

Example 5. Preparation of the quinoline ligand (Formula 7).

Formula 7

5 A mixture of 2-chloromethyl-8-hydroxyquinoline hydrochloride (Chem-Master International) (0.69 g, 3 mmol), ethylenediamine- N,N,N' -triacetic acid tri(*t*-butyl)ester (10.8 g, 3 mmol), diisopropylethylamine (0.78 g, 6 mmol), and sodium iodide (0.15 g, 1 mmol) in dimethoxyethane (10 mL) was heated under reflux for 4 hours. The reaction mixture was poured onto

10 water and extracted with ether (3 x 30 mL). The combined organic layers were washed with water (3 x 50 mL), dried (Na_2SO_4), filtered, and the filtrate evaporated in vacuo to furnish the triester which was purified by silica gel chromatography using chloroform-methanol (9:1) as eluent. $^1\text{H-NMR}$ (CDCl_3) δ 8.10 (d, 1H), 7.61 (d, 1H), 7.32 (m, 2H), 7.08 (d, 1H), 4.08 (s, 2H),

15 3.42 (s, 4H), 3.38 (s, 4H), 2.83 (bs, 4H); $^{13}\text{C-NMR}$ (CDCl_3) δ 170.6, 158.0, 152.1, 137.4, 136.3, 127.5, 127.1, 122.1, 117.5, 112.8, 109.9, 80.9, 80.8, 60.4, 56.1, 55.8, 52.5, 51.8.


The triester obtained above (820 mg) was treated with 96% formic acid (10 mL) and heated at 80-90 °C for 15 minutes and thereafter kept at ambient

20 temperature for 24 hours. Excess formic acid was removed by evaporation in vacuo and the residue was triturated with acetone (50 mL). The solid was collected by filtration and dried to give 520 mg of the ligand of Formula 7 as a pale pink solid. $^1\text{H-NMR}$ (DMSO-d_6) δ 8.15 (d, 1H), 7.60 (d, 1H), 7.32 (m, 2H), 7.05 (d, 1H), 4.25 (s, 2H), 3.42 (s, 4H), 3.38 (s, 4H), 2.80 (bs, 4H); $^{13}\text{C-NMR}$ (DMSO-d_6) δ 172.4, 171.3, 156.2, 152.8, 137.3, 136.6, 127.8, 127.2,

25


121.4, 117.6, 111.2, 59.3, 55.0, 54.8, 51.7, 50.9; electrospray mass spectrum, $m/Z = 392$ ($M + H$).

Example 6. Preparation of europium complex (Formula 8).




Formula 8

10 A mixture of the ligand of Formula 7 (780 mg, 2 mmol) and europium oxide (352 mg, 1 mmol) in deionized, distilled water (10 mL) was heated under reflux for 72 hours. The mixture remained heterogeneous throughout the heating period. The precipitate was filtered, washed with water and dried to give the europium complex of Formula 8 as an off-white solid.

Example 7. Hypothetical preparation of lutetium complex (Formula 9).

Formula 9


A mixture of the ligand of Formula 7 (798 mg, 2 mmol) and lutetium
 5 oxide (398 mg, 1 mmol) in deionized, distilled water (10 mL) is heated under
 reflux for 24 hours. The solution is filtered through fine porosity sintered
 glass funnel to remove undissolved impurities and the filtrate is poured onto
 acetone (200 mL). The precipitate is collected, washed with acetone, and
 dried. The crude lutetium complex of Formula 9 is purified by C-18 reverse
 10 phase chromatography.

Example 8. Preparation of iron complex (Formula 10).

Formula 10

A solution of the ligand of Formula 7 (391 mg, 1 mmol) in 1N sodium
 15 hydroxide was treated with ferric chloride hexahydrate (269 mg, 1 mmol) and
 the entire mixture was stirred at ambient temperature for 16 hours. The dark
 reaction mixture was filtered and the filtrate washed with ice-cold water and
 dried. The U.V. spectrum showed intense bands at 455 nm and 596 nm. The
 20 mass spectrum showed the correct molecular ion for the iron complex of
 Formula 10, $m/Z = 445 (M + H)$.


Example 9. Hypothetical preparation of the gadolinium complex (Formula 11).



Formula 11

A mixture of the ligand of Formula 7 (780 mg, 2 mmol) and gadolinium oxide (362 mg, 1 mmol) in deionized, distilled water (10 mL) is heated under reflux for 24 hours. The solution is filtered through fine porosity sintered glass funnel to remove undissolved impurities and the filtrate is poured onto acetone (200 mL). The precipitate is collected, washed with acetone, and dried. The crude gadolinium complex of Formula 11 is purified by C-18 reverse phase chromatography.

Example 10. Preparation of the quinoline ligand (Formula 12).




Formula 12

A mixture of 2-chloromethyl-8-hydroxyquinoline hydrochloride (1.05 g, 5 mmol), di-*t*-butyl 3,10-bis(*t*-butoxycarbonylmethyl)-3,6,10-triazadodecanedioate (1.78 g, 1 mmol), diisopropylethylamine (3.87 g, 20

mmol), and sodium iodide (0.15 g, 1 mmol) in dimethoxyethane (10 mL) was heated under reflux for 4 hours. The reaction mixture was poured onto water and extracted with ethyl ether (3 x 20 mL). The combined organic layers were washed with water (3 x 20 mL), dried (Na_2SO_4), filtered, and the filtrate
5 evaporated in vacuo to furnish the tetraester which was purified by silica gel chromatography using chloroform-methanol (9:1). The yield of the pale brown tetraester was 1.8 g. $^1\text{H-NMR}$ (CDCl_3) δ 8.02 (d, 1H), 7.61 (d, 1H), 7.38 (m, 2H), 7.25 (d, 1H), 7.15 (d, 1H), 3.95 (s, 2H), 3.42 (bs, 8H), 2.90 (m, 4H), 2.72 (bs, 4H), 1.21 (s, 48H); $^{13}\text{C-NMR}$ (CDCl_3) δ 170.6, 158.7, 152.1,
10 137.4, 136.2, 127.5, 127.0, 122.0, 117.5, 112.8, 109.9, 80.8, 60.8, 56.1, 53.0, 51.8, 28.1.


The tetraester obtained above was treated with 96% formic acid (10 mL) and kept at ambient temperature for 24 hours. The reaction mixture was poured onto acetone (200 mL) and solid was collected by filtration and dried
15 to furnish the ligand of Formula 12 as an off-white solid. $^1\text{H-NMR}$ (DMSO-d_6) δ 8.32 (d, 1H), 7.58 (d, 1H), 7.38 (m, 2H), 7.05 (d, 1H), 4.58 (bs, 2H), 3.37 (bs, 8H), 3.12 (bs, 4H), 3.00 (bs, 4H); $^{13}\text{C-NMR}$ (DMSO-d_6) δ 173.0, 153.1, 152.4, 137.2, 136.6, 137.0, 127.8, 127.7, 121.3, 117.4, 111.5, 58.0, 55.6, 51.7, 49.7; electrospray mass spectrum, $m/z = 493$ ($M + H$).

20

Example 11. Preparation of the europium complex (Formula 13).

Formula 13

- A mixture of the ligand of Formula 12 (984 mg, 2 mmol) and
 5 europium oxide (352 mg, 1 mmol) in deionized, distilled water (10 mL) was
 heated under reflux for 24 hours. The solution was filtered through a fine
 porosity sintered glass funnel to remove undissolved impurities and the
 filtrate was poured onto acetone (200 mL). The yellow-orange solid
 precipitate, a crude europium complex of Formula 13, was collected, washed
 10 with acetone, and dried.


Example 12. Hypothetical preparation of the lutetium complex (Formula 14).

Formula 14

- A mixture of the ligand of Formula 12 (984 mg, 2 mmol) and lutetium
 15 oxide (398 mg, 1 mmol) in deionized, distilled water (10 mL) is heated under
 reflux for 24 hours. The solution is filtered through fine porosity sintered
 glass funnel to remove undissolved impurities and the filtrate is poured onto


acetone (200 mL). The precipitate is collected, washed with acetone, and dried. The crude lutetium complex of Formula 14 is purified by C-18 reverse phase chromatography.

5 Example 13. Hypothetical preparation of the chromium complex (Formula 15).




Formula 15

A mixture of the ligand of Formula 12 (390 mg, 1 mmol) and
10 chromium acetylacetonate (350 mg, 1 mmol) in dimethylformamide (3 mL)
is treated with two drops of water and the entire mixture was heated at 100-
120 °C for 24 hours. After cooling the reaction mixture to ambient
temperature, the solution is poured onto ethyl ether. The precipitate is
collected by filtration, washed with ether, and dried. The crude chromium
15 complex of Formula 15 is purified by crystallization or C-18 reverse phase
chromatography.

Example 14. Hypothetical preparation of gadolinium complex (Formula 16).

Formula 16

- A mixture of the ligand of Formula 12 (780 mg, 2 mmol) and
 5 gadolinium oxide (362 mg, 1 mmol) in deionized, distilled water (10 mL) is heated under reflux for 24 hours. The solution is filtered through a fine porosity sintered glass funnel to remove undissolved impurities and the filtrate is poured onto acetone (200 mL). The precipitate is collected, washed with acetone, and dried. The crude gadolinium complex of Formula 16 is
 10 purified by C-18 reverse phase chromatography.

Example 15. Preparation of the quinoline ligand (Formula 17).


Formula 17

- A mixture of 2-aminomethyl-8-hydroxyquinoline hydrochloride (1.05 g, 5 mmol) N-succinimidyl S-tetrahydropyranylmethylmercaptoacetate (1.37 g, 5 mmol), and triethylamine (0.51 g, 5 mmol) in acetonitrile (15 mL) was heated under reflux for 2 hours. The reaction mixture was poured onto water and extracted with methylene chloride (3 x 20 mL). The combined organic layers
 20 were washed with water and dried (MgSO₄) to give the ligand of Formula 17

as an orange gum which was sufficiently pure for complexation purposes.

$^1\text{H-NMR}$ (CDCl_3) δ 8.42 (broad, 1H), 8.20 (d, 1H), 7.41 (d, 1H), 7.35 (m, 2H), 7.18 (d, 1H), 4.95 (dd, 1H), 4.82 (dd, 1H), 4.73 (dd, 1H), 4.00 (m, 1H), 3.78 (m, 1H), 3.57 (d, 1H), 3.35 (d, 1H), 2.10-1.55 (m, 6H); $^{13}\text{C-NMR}$ (CDCl_3) δ 169.6, 154.1, 152.1, 137.0, 136.9, 127.5, 127.4, 120.2, 117.8, 110.9, 83.7, 65.5, 44.7, 35.2, 30.8, 25.0, 21.6.

Example 16. Preparation of rhenium complex (Formula 18).




Formula 18

A mixture of the ligand of Formula 17 (332 mg, 1 mmol), triethylamine (2 drops), and oxodichlorobis(triphenyl)phosphinoethoxyrhenium (880 mg, 1 mmol) in ethanol (5 ml) was heated under reflux for 6 hours. The precipitate was collected by filtration and the crude rhenium complex of Formula 18 was purified by recrystallization.

While the invention has been disclosed by reference to the details of preferred embodiments of the invention, it is to be understood that the disclosure is intended in an illustrative rather than in a limiting sense, as it is contemplated that modifications will readily occur to those skilled in the art, within the spirit of the invention and the scope of the appended claims.

We claim:

1. A composition comprising a quinoline ligand of Formula I:



Formula I

5

- wherein R¹ to R⁵ may be the same or different and are selected from the group consisting of hydrogen, C1-C10 alkyl, -OH, C1-C10 polyhydroxyalkyl, C1-C10 alkoxy, C1-C10 alkoxyalkyl, -SO₃H, -(CH₂)_m-CO₂H, and -NR⁶R⁷; R⁶ and R⁷ may be the same or different and are selected from the group consisting of hydrogen, C1-C10 alkyl, C1-C10 aryl, and C1-C10 polyhydroxyalkyl; m ranges from 0 to 10; A¹ is selected from the group consisting of -OH, -CO₂H, -N(R⁸)(R⁹), -SPg, -CONHR¹⁰, and -HNCOR¹¹; R⁸ and R⁹ may be the same or different and are selected from the group consisting of hydrogen, C1-C10 alkyl, C1-C10 aryl, C1-C10 polyhydroxyalkyl, -(CH₂)_m-CO₂H, and -(CH₂)₂-N(CH₂CO₂H)₂; R¹⁰ is selected from the group consisting of hydrogen, C1-C10 alkyl, C1-C10 aryl, C1-C10 polyhydroxyalkyl, and -(CH₂)₂-SPg; R¹¹ is selected from the group consisting of hydrogen, C1-C10 alkyl, C1-C10 aryl, C1-C10 polyhydroxyalkyl, and -CH(R¹²)-SPg; R¹² is selected from the group consisting of hydrogen, C1-C10 alkyl, C1-C10 aryl, C1-C10 polyhydroxyalkyl, -(CH₂)_m-CO₂H and -(CH₂)_n-NH₂; n varies from 1 to 10; B¹ is selected from the group consisting of -CHR¹³ and -CH(R¹⁴)(R¹⁵); R¹³ to R¹⁵ may be the same or different and are defined in the same manner as R¹²; C¹ is selected from the group consisting of hydrogen, C1-C10 alkyl, C1-C10 aryl, C1-C10 polyhydroxyalkyl, C1-C10 alkoxy, C1-C10 alkoxyalkyl, -
- 10
- 15
- 20

(CH₂)_mCO₂H, -CH₂CH₂-N(CH₂CO₂H)₂, -(CH₂)₂-SPg, and -COCH(R¹⁶)-SPg; R¹⁶ is defined in the same manner as R¹²; D¹ is selected from the group consisting of hydrogen, C1-C10 alkyl, C1-C10 aryl, hydroxyl, C1-C10 polyhydroxyalkyl, C1-C10 alkoxy, C1-C10 alkoxyalkyl, -(CH₂)_mCO₂H, and
 5 -CH₂CH₂-N(CH₂CO₂H)₂; and Pg is hydrogen or a protecting group selected from the group consisting of t-butyl, benzyl, benzoyl, 2,4-dimethoxybenzyl, trityl, tetrahydropyranyl, C1-C10 acyl, C1-C10 alkoxyalkyl, and C1-C10 alkoxycarbonyl.


2. The composition according to Claim 1 wherein R¹ to R⁵ are selected
 10 from the group consisting of hydrogen, -OH, C1-C10 alkoxy, -(CH₂)_m-CO₂H, and -N(R⁶)(R⁷); A¹ is selected from the group consisting of -OH, -N(R⁸)(R⁹), and -HNCOR¹¹; B¹ is -CHR¹³; C¹ is selected from the group consisting of hydrogen, C1-C10 alkyl, -(CH₂)_mCO₂H, -CH₂CH₂-N(CH₂CO₂H)₂, and -COCH(R¹⁶)-SPg; D¹ is selected from the group
 15 consisting of hydrogen, C1-C10 alkyl, C1-C10 polyhydroxyalkyl, -(CH₂)_m-CO₂H, and -CH₂CH₂-N(CH₂CO₂H)₂; and Pg is selected from the group consisting of C1-C10 acyl, tetrahydropyranyl, C1-C10 alkoxyalkyl, and C1-C10 alkoxycarbonyl.

3. The composition according to Claim 2 wherein R¹ to R⁵ are hydrogens;
 20 A¹ is -OH or -N(CH₂CO₂H)₂; B¹ is -CH₂-; C¹ is selected from the group consisting of -CH₂-CO₂H, -CH₂CH₂-N(CH₂CO₂H)₂, and -COCH₂-SPg; D¹ is selected from the group consisting of hydrogen, C1-C10 alkyl, and -CH₂-CO₂H; and Pg is selected from the group consisting of benzoyl, tetrahydropyranyl, and methoxycarbonyl.

4. The composition according to Claim 3 wherein R¹ to R⁵ are hydrogens;
 25 A¹ is -OH; B¹ is -CH₂-; C¹ is -CH₂-CO₂H; and D¹ is -CH₂-CO₂H.

5. The composition according to Claim 3 wherein R¹ to R⁵ are hydrogens;
 A¹ is -OH; B¹ is -CH₂-; C¹ is -COCH₂-SPg; D¹ is hydrogen; and Pg is a tetrahydropyranyl group.

6. The composition according to Claim 3 wherein R^1 to R^5 are hydrogens; A^1 is $-OH$; B^1 is $-CH_2-$; C^1 is $-CH_2CH_2-N(CH_2CO_2H)_2$; and D^1 is $-CH_2-CO_2H$.
7. The composition according to Claim 3 wherein R^1 to R^5 are hydrogens; A^1 is $-OH$; B^1 is $-CH_2-$; and C^1 and D^1 are $-CH_2CH_2-N(CH_2CO_2H)_2$.
8. The composition according to Claim 3 wherein R^1 to R^5 are hydrogens; A^1 is $-N(CH_2CO_2H)_2$; B^1 is $-CH_2-$; C^1 is $-CH_2CO_2H$; and D^1 is $-CH_2-CO_2H$.
9. A composition comprising a quinoline metal complex of Formula II:



Formula II

10

wherein R^{17} to R^{21} may be the same or different and are selected from the group consisting of hydrogen, C1-C10 alkyl, $-OH$, C1-C10 polyhydroxyalkyl, C1-C10 alkoxy, C1-C10 alkoxyalkyl, $-SO_3H$, $-(CH_2)_m-CO_2H$, and $-NR^6R^7$; A^2 is selected from the group consisting of $-O-$, $-CO_2-$, $-N(R^8)(R^9)$, $-SPg-$

15

$CON(R^{10})$, and $-NCOR^{11}$; R^8 and R^9 may be the same or different and are selected from the group consisting of hydrogen, C1-C10 alkyl, C1-C10 aryl, C1-C10 polyhydroxyalkyl, $-(CH_2)_mCO_2-$, and $-(CH_2)_2-N(CH_2CO_2-)_2$; R^{10} is selected from the group consisting of hydrogen, C1-C10 alkyl, C1-C10 aryl, C1-C10 polyhydroxyalkyl, and $-(CH_2)_2-S-$; R^{11} is selected from the group

20

consisting of hydrogen, C1-C10 alkyl, C1-C10 aryl, C1-C10 polyhydroxyalkyl, and $-CH(R^{12})-S-$; R^{12} is selected from the group consisting of hydrogen, C1-C10 alkyl, C1-C10 aryl, C1-C10 polyhydroxyalkyl, $-(CH_2)_mCO_2-$ and $-(CH_2)_nNH_2$; n varies from 1 to 10; B^2 is selected from the group consisting of $-CHR^{13}$ and $-CH(R^{14})CH(R^{15})$; R^{13} to R^{15} may be the same


or different and are selected from the group consisting of hydrogen, C1-C10 alkyl, C1-C10 aryl, C1-C10 polyhydroxyalkyl, $-(CH_2)_mCO_2^-$ and $-(CH_2)_nNH_2$; C² is selected from the group consisting of hydrogen, C1-C10 alkyl, C1-C10 aryl, C1-C10 polyhydroxyalkyl, C1-C10 alkoxy, C1-C10 alkoxyalkyl, -
 5 $(CH_2)_mCO_2^-$, $-CH_2CH_2-N(CH_2CO_2^-)_2$, $-(CH_2)_2-S^-$, and $-COCH(R^{16})-S^-$; R¹⁶ is selected from the group consisting of hydrogen, C1-C10 alkyl, C1-C10 aryl, C1-C10 polyhydroxyalkyl, $-(CH_2)_mCO_2^-$ and $-(CH_2)_nNH_2$; D² is selected from the group consisting of hydrogen, C1-C10 alkyl, C1-C10 aryl, hydroxyl, C1-C10 polyhydroxyalkyl, C1-C10 alkoxy, C1-C10 alkoxyalkyl, $-(CH_2)_mCO_2^-$,
 10 and $-CH_2CH_2-N(CH_2CO_2^-)_2$; and M is a metal ion having an atomic number of 24 to 31, 42 to 49, 58-71, or 74-83.

10. The composition according to Claim 9 wherein R¹⁷ to R²¹ are selected from the group consisting of hydrogen, -O⁻, C1-C10 alkoxy, $-(CH_2)_m-CO_2^-$, and $-N(R^6)(R^7)$; A² is selected from the group consisting of -O⁻, $-N(R^8)(R^9)$,
 15 and $-HNCOR^{11}$; B² is $-CHR^{13}$; C² is selected from the group consisting of hydrogen, C1-C10 alkyl, $-(CH_2)_mCO_2^-$, $-CH_2CH_2-N(CH_2CO_2^-)_2$, and $-COCH(R^{16})-S^-$; D² is selected from the group consisting of hydrogen, C1-C10 alkyl, C1-C10 polyhydroxyalkyl, $-(CH_2)_m-CO_2^-$, and $-CH_2CH_2-N(CH_2CO_2^-)_2$; and M is a metal ion having an atomic number of 24 to 28, 31,
 20 42 to 45, 49, 62-65, 71, 75, 76, or 83.

11. The composition according to Claim 9 wherein R¹⁷ to R²¹ are hydrogens; A² is -O⁻ or $-N(CH_2CO_2^-)_2$; B² is $-CH_2-$; C² is selected from the group consisting of $-CH_2-CO_2^-$, $-CH_2CH_2-N(CH_2CO_2^-)_2$, and $-COCH_2-S^-$; D² is selected from the group consisting of hydrogen, C1-C10 alkyl, and $-CH_2-$
 25 CO_2^- ; and M is a metal ion having an atomic number of 24-26, 28, 31, 43, 44, 49, 62-65, 71, 75, or 76.

12. The composition according to Claim 11 wherein R¹⁷ to R²¹ are hydrogens; A² is -O⁻; B² is $-CH_2-$; C² is $-CH_2-CO_2^-$; D² is $-CH_2-CO_2^-$; and M is a metal ion having an atomic number of 24-26, 31, 44, 49, or 63.

13. The composition according to Claim 11 wherein R^{17} to R^{21} are hydrogens; A^2 is $-O^-$; B^2 is $-CH_2-$; C^2 is $-COCH_2-S-$; D^2 is hydrogen; and M is a metal ion having an atomic number of 43, 44, 75, or 76.
14. The composition according to Claim 11 wherein R^{17} to R^{21} are hydrogens; A^2 is $-O^-$; B^2 is $-CH_2-$; C^2 is $-CH_2CH_2-N(CH_2CO_2^-)_2$; D^2 is $-CH_2-CO_2^-$; and M is a metal ion having an atomic number of 24-26, 44, 49, 62-65, or 71.
15. The composition according to Claim 10 wherein R^{17} to R^{21} are hydrogens; A^2 is $-O^-$; B^2 is $-CH_2-$; C^2 and D^2 are $-CH_2CH_2-N(CH_2CO_2^-)_2$; and M is a metal ion having an atomic number of 24-26, 44, 49, 62-65, or 71.
16. The composition according to Claim 11 wherein R^{17} to R^{21} are hydrogens; A^2 is $-N(CH_2CO_2^-)_2$; B^2 is $-CH_2-$; C^2 is $-CH_2CO_2^-$; D^2 is $-CH_2-CO_2^-$; and M is a metal ion having an atomic number of 24-26, 44, 49, 62-65, or 71.
17. A method of performing a diagnostic or a therapeutic procedure
- 15 comprising administering an effective amount of the quinoline metal complex of Formula II:



Formula II

- 20 wherein R^{17} to R^{21} may be the same or different and are selected from the group consisting of hydrogen, C1-C10 alkyl, $-OH$, C1-C10 polyhydroxyalkyl, C1-C10 alkoxy, C1-C10 alkoxyalkyl, $-SO_3H$, $-(CH_2)_m-CO_2H$, and $-NR^6R^7$; A^2 is selected from the group consisting of $-O^-$, $-CO_2^-$, $-N(R^8)(R^9)$, $-SPg-$, $CON(R^{10})$, and $-NCOR^{11}$; R^8 and R^9 may be the same or different and are

- selected from the group consisting of hydrogen, C1-C10 alkyl, C1-C10 aryl, C1-C10 polyhydroxyalkyl, $-(CH_2)_mCO_2^-$, and $-(CH_2)_2-N(CH_2CO_2^-)_2$; R^{10} is selected from the group consisting of hydrogen, C1-C10 alkyl, C1-C10 aryl, C1-C10 polyhydroxyalkyl, and $-(CH_2)_2-S^-$; R^{11} is selected from the group consisting of hydrogen, C1-C10 alkyl, C1-C10 aryl, C1-C10 polyhydroxyalkyl, and $-CH(R^{12})-S^-$; R^{12} is selected from the group consisting of hydrogen, C1-C10 alkyl, C1-C10 aryl, C1-C10 polyhydroxyalkyl, $-(CH_2)_mCO_2^-$ and $-(CH_2)_nNH_2$; n varies from 1 to 10; B^2 is selected from the group consisting of $-CHR^{13}$, and $-CH(R^{14})CH(R^{15})$; R^{13} to R^{15} may be the same or different and are selected from the group consisting of hydrogen, C1-C10 alkyl, C1-C10 aryl, C1-C10 polyhydroxyalkyl, $-(CH_2)_mCO_2^-$ and $-(CH_2)_nNH_2$; C^2 is selected from the group consisting of hydrogen, C1-C10 alkyl, C1-C10 aryl, C1-C10 polyhydroxyalkyl, C1-C10 alkoxy, C1-C10 alkoxyalkyl, $-(CH_2)_mCO_2^-$, $-CH_2CH_2-N(CH_2CO_2^-)_2$, $-(CH_2)_2-S^-$, and $-COCH(R^{16})-S^-$; R^{16} is selected from the group consisting of hydrogen, C1-C10 alkyl, C1-C10 aryl, C1-C10 polyhydroxyalkyl, $-(CH_2)_mCO_2^-$ and $-(CH_2)_nNH_2$; D^2 is selected from the group consisting of hydrogen, C1-C10 alkyl, C1-C10 aryl, hydroxyl, C1-C10 polyhydroxyalkyl, C1-C10 alkoxy, C1-C10 alkoxyalkyl, $-(CH_2)_mCO_2^-$, and $-CH_2CH_2-N(CH_2CO_2^-)_2$; and M is a metal ion having an atomic number of 24-31, 42-49, 58-71, or 74-83.
18. The method according to Claim 17 wherein R^{17} to R^{21} are selected from the group consisting of hydrogen, $-O^-$, C1-C10 alkoxy, $-(CH_2)_mCO_2^-$, and $-N(R^6)(R^7)$; A^2 is selected from the group consisting of $-O^-$, $-N(R^8)(R^9)$, and $-HNCOR^{11}$; B^2 is $-CHR^{13}$; C^2 is selected from the group consisting of hydrogen, C1-C10 alkyl, $-(CH_2)_mCO_2^-$, $-CH_2CH_2-N(CH_2CO_2^-)_2$, and $-COCH(R^{16})-S^-$; D^2 is selected from the group consisting of hydrogen, C1-C10 alkyl, C1-C10 polyhydroxyalkyl, $-(CH_2)_mCO_2^-$, and $-CH_2CH_2-N(CH_2CO_2^-)_2$; and M is a metal ion having an atomic number of 24-28, 31, 42-45, 49, 62-65, 71, 75, 76, or 83.

19. The method according to Claim 17 wherein R^{17} to R^{21} are hydrogens; A^2 is $-O^-$ or $-N(CH_2CO_2^-)_2$; B^2 is $-CH_2^-$; C^2 is selected from the group consisting of $-CH_2-CO_2^-$, $-CH_2CH_2-N(CH_2CO_2^-)_2$, and $-COCH_2-S^-$; D^2 is selected from the group consisting of hydrogen, C1-C10 alkyl, and $-CH_2-CO_2^-$; and M is a metal ion having an atomic number of 24-26, 28, 31, 43, 44, 49, 62-65, 71, 75, or 76.
20. The method according to Claim 19 wherein R^{17} to R^{21} are hydrogens; A^2 is $-O^-$; B^2 is $-CH_2^-$; C^2 is $-CH_2-CO_2^-$; D^2 is $-CH_2-CO_2^-$; and M is a metal ion having an atomic number of 24-26, 31, 44, 49, or 63.
- 10 21. The method according to Claim 19 wherein R^{17} to R^{21} are hydrogens; A^2 is $-O^-$; B^2 is $-CH_2^-$; C^2 is $-COCH_2-S^-$; D^2 is hydrogen; and M is a metal ion having an atomic number of 43, 44, 75, or 76.
22. The method according to Claim 19 wherein R^{17} to R^{21} are hydrogens; A^2 is $-O^-$; B^2 is $-CH_2^-$; C^2 is $-CH_2CH_2-N(CH_2CO_2^-)_2$; D^2 is $-CH_2-CO_2^-$; and M is a metal ion having an atomic number of 24-26, 44, 49, 62-65, or 71.
- 15 23. The method according to Claim 18 wherein R^{17} to R^{21} are hydrogens; A^2 is $-O^-$; B^2 is $-CH_2^-$; C^2 and D^2 are $-CH_2CH_2-N(CH_2CO_2^-)_2$; and M is a metal ion having an atomic number of 24-26, 44, 49, 62-65, or 71.
24. The method according to Claim 19 wherein R^{17} to R^{21} are hydrogens;
- 20 A^2 is $-N(CH_2CO_2^-)_2$; B^2 is $-CH_2^-$; C^2 is $-CH_2CO_2^-$; D^2 is $-CH_2-CO_2^-$; and M is a metal ion having an atomic number of 24-26, 44, 49, 62-65, or 71.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US01/06394

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : C07D 265/30; A61K 31/555; C07F 5/02
US CL : 544/176; 546/7; 534/16; 514/185

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
U.S. : 544/176; 546/7; 534/16; 514/185

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
CAS online structure search

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5,118,499 A1 (THEODOROPULOS et al) 02 June 1992, column 2.	1-9
X	WO 90/07376 A1 (EXPLOATERINGS AB T.B.F.) 12 July 1990, se entire document in particular example 9 on page 22.	1-9
X	Chem. Abstr. Vol. 87, 1977. MUKHOPADHY et al "Synthesis of possible antiamebic agents". J. Heterocycles. Abstract No. 87:39245, RN63218-48-4, J. Indian Chem. Soc. (1976) 53(10), 1838-40.	1-9

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T"

later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X"

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y"

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"Z"

document member of the same patent family

Date of the actual completion of the international search

25 April 2001 (25.04.2001)

Date of mailing of the international search report

05 JUN 2001

Name and mailing address of the ISA/US

Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231
Facsimile No. (703)305-3230

Authorized officer

Mukund Shah

Telephone No. 703 388-1235

JOYCE BRIDGERS
PARALEGAL SPECIALIST
CHEMICAL MATRIX